

**PSMD4 antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI10048****Specification**

---

**PSMD4 antibody - C-terminal region - Product Information**

Application	IHC, WB
Primary Accession	<a href="#">P55036</a>
Other Accession	<a href="#">P55036</a> , <a href="#">NP_002801</a> , <a href="#">NM_002810</a>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Sheep, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Pig, Sheep, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41 kDa KDa

**PSMD4 antibody - C-terminal region - Additional Information****Gene ID 5710**

Alias Symbol **AF, ASF, S5A, AF-1, MCB1, Rpn10, pUB-R5**

**Other Names**

26S proteasome non-ATPase regulatory subunit 4, 26S proteasome regulatory subunit RPN10, 26S proteasome regulatory subunit S5A, Antisecretory factor 1, AF, ASF, Multiubiquitin chain-binding protein, PSMD4, MCB1

**Target/Specificity**

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits, 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMD4 encodes one of the non-ATPase subunits of the 19S regulator lid.

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-PSMD4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

PSMD4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## PSMD4 antibody - C-terminal region - Protein Information

**Name** PSMD4

**Synonyms** MCB1

### Function


Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair. PSMD4 acts as an ubiquitin receptor subunit through ubiquitin- interacting motifs and selects ubiquitin-conjugates for destruction. Displays a preferred selectivity for longer polyubiquitin chains.

## PSMD4 antibody - C-terminal region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## PSMD4 antibody - C-terminal region - Images



Human Brain  
Rabbit Anti-PSMD4 Antibody  
Paraffin Embedded Tissue: Human neural cell  
Cellular Data: Epithelial cells of renal tubule  
Antibody Concentration: 4.0-8.0 µg/ml  
Magnification: 400X



90 kDa  
60 kDa  
42 kDa  
32 kDa  
23 kDa

PSMD4 antibody - C-terminal region (AI10048) in Human Raji cells using Western Blot

WB Suggested Anti-PSMD4 Antibody Titration: 0.2-1  $\mu$ g/ml

Positive Control: Raji cell lysate

PSMD4 is strongly supported by BioGPS gene expression data to be expressed in Human Raji cells

#### **PSMD4 antibody - C-terminal region - Background**

This is a rabbit polyclonal antibody against PSMD4. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire ([sales@abgent.com](mailto:sales@abgent.com)).