

## RUNX1T1 antibody - middle region

Rabbit Polyclonal Antibody Catalog # Al10108

# **Specification**

## **RUNX1T1** antibody - middle region - Product Information

Application WB
Primary Accession Q06455

Other Accession <u>Q06455, NP 783554, NM 175636</u>

Reactivity Human, Mouse, Rat, Rabbit, Zebrafish, Pig,

Dog, Guinea Pig, Horse, Bovine

Predicted Human, Mouse, Rat, Zebrafish, Pig,

Chicken, Dog, Horse

Host Rabbit
Clonality Polyclonal
Calculated MW 62 kDa KDa

# **RUNX1T1** antibody - middle region - Additional Information

Gene ID 862

Alias Symbol CDR, ETO, MTG8, AML1T1, ZMYND2, CBFA2T1

#### **Other Names**

Protein CBFA2T1, Cyclin-D-related protein, Eight twenty one protein, Protein ETO, Protein MTG8, Zinc finger MYND domain-containing protein 2, RUNX1T1, AML1T1, CBFA2T1, CDR, ETO, MTG8, ZMYND2

### Target/Specificity

RUNX1T1 is a putative zinc finger transcription factor and oncoprotein. In acute myeloid leukemia, especially in the M2 subtype, the t(8, 21)(q22, q22) translocation is one of the most frequent karyotypic abnormalities. The translocation produces a chimeric gene made up of the 5'-region of the RUNX1 gene fused to the 3'-region of this gene. The chimeric protein is thought to associate with the nuclear corepressor/histone deacetylase complex to block hematopoietic differentiation.

#### **Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

#### **Reconstitution & Storage**

Add 100 ul of distilled water. Final anti-RUNX1T1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

## **Precautions**

RUNX1T1 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

#### **RUNX1T1** antibody - middle region - Protein Information



#### Name RUNX1T1

#### Synonyms AML1T1, CBFA2T1, CDR, ETO, MTG8, ZMYND2

#### **Function**

Transcriptional corepressor which facilitates transcriptional repression via its association with DNA-binding transcription factors and recruitment of other corepressors and histone-modifying enzymes (PubMed:<a href="http://www.uniprot.org/citations/12559562" http://www.uniprot.org/citations/12559562" http://www.uniprot.org/citations/12559562"

target="\_blank">12559562</a>, PubMed:<a href="http://www.uniprot.org/citations/15203199" target="\_blank">15203199</a>, PubMed:<a href="http://www.uniprot.org/citations/10688654" target="\_blank">10688654</a>). Can repress the expression of MMP7 in a ZBTB33-dependent manner (PubMed:<a href="http://www.uniprot.org/citations/23251453"

target="\_blank">23251453</a>). Can repress transactivation mediated by TCF12 (PubMed:<a href="http://www.uniprot.org/citations/16803958" target="\_blank">16803958</a>). Acts as a negative regulator of adipogenesis (By similarity). The AML1-MTG8/ETO fusion protein frequently found in leukemic cells is involved in leukemogenesis and contributes to hematopoietic stem/progenitor cell self-renewal (PubMed:<a href="http://www.uniprot.org/citations/23812588" target="\_blank">23812588</a>).

#### **Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00440, ECO:0000269|PubMed:10973986}. Note=Colocalizes with ATN1 in discrete nuclear dots

#### **Tissue Location**


Most abundantly expressed in brain. Lower levels in lung, heart, testis and ovary

### RUNX1T1 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### **RUNX1T1** antibody - middle region - Images




RUNX1T1 antibody - middle region (Al10108) in Human 721\_B cells using Western Blot Host: Rabbit



Target Name: WT1 Sample Tissue: 721 B

Antibody Dilution: 1.0µg/mlRUNX1T1 is supported by BioGPS gene expression data to be

expressed in 721\_B



RUNX1T1 antibody - middle region (Al10108) in Human HepG2 cells using Western Blot

WB Suggested Anti-RUNX1T1 Antibody Titration: 1.25µg/ml

Positive Control: HepG2 cell lysate

There is BioGPS gene expression data showing that RUNX1T1 is expressed in HepG2

# RUNX1T1 antibody - middle region - Background

This is a rabbit polyclonal antibody against RUNX1T1. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).