

FHL2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10132**Specification**

FHL2 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q14192
Other Accession	Q14192 , NP_963849 , NM_201555
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Chicken, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32 kDa KDa

FHL2 antibody - C-terminal region - Additional Information**Gene ID 2274**

Alias Symbol **DRAL, AAG11, FHL-2, SLIM3, SLIM-3**
Other Names
Four and a half LIM domains protein 2, FHL-2, LIM domain protein DRAL, Skeletal muscle LIM-protein 3, SLIM-3, FHL2, DRAL, SLIM3

Target/Specificity

FHL2 is a member of LIM proteins that contain a highly conserved double zinc finger motif called the LIM domain.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-FHL2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

FHL2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

FHL2 antibody - C-terminal region - Protein Information

Name FHL2

Synonyms DRAL, SLIM3

Function

May function as a molecular transmitter linking various signaling pathways to transcriptional regulation. Negatively regulates the transcriptional repressor E4F1 and may function in cell growth. Inhibits the transcriptional activity of FOXO1 and its apoptotic function by enhancing the interaction of FOXO1 with SIRT1 and FOXO1 deacetylation. Negatively regulates the calcineurin/NFAT signaling pathway in cardiomyocytes (PubMed:28717008).

Cellular Location

Cytoplasm. Nucleus. Cytoplasm, myofibril, sarcomere, Z line {ECO:0000250|UniProtKB:O35115}

Tissue Location

Expressed in skeletal muscle and heart.


FHL2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FHL2 antibody - C-terminal region - Images

FHL2 antibody - C-terminal region (AI10132) in Human Heart cells using Immunohistochemistry
Human Heart

A Western blot analysis showing a single protein band. The molecular weight markers on the left are 90 kDa, 60 kDa, 42 kDa, 32 kDa, and 23 kDa. A distinct band is visible at approximately 32 kDa.

90 kDa__
60 kDa__
42 kDa__
32 kDa__
23 kDa__

FHL2 antibody - C-terminal region (AI10132) in Human heart cells using Western Blot
WB Suggested Anti-FHL2 Antibody Titration: 0.6µg/ml
ELISA Titer: 1:312500
Positive Control: Human heart

FHL2 antibody - C-terminal region - Background

This is a rabbit polyclonal antibody against FHL2. It was validated on Western Blot and immunohistochemistry by Abgent. At Abgent we manufacture rabbit polyclonal antibodies on a large scale (200-1000 products/month) of high throughput manner. Our antibodies are peptide based and protein family oriented. We usually provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).