

UHRF2 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10140**Specification**

UHRF2 antibody - middle region - Product Information

Application	WB
Primary Accession	O96PU4
Other Accession	O96PU4 , NP_690856 , NM_152896
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Chicken, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	90 kDa KDa

UHRF2 antibody - middle region - Additional Information**Gene ID** 115426**Alias Symbol** DKFZp434B0920, DKFZp686G0837, MGC33463, NIRF, RNF107, RP11-472F14.2, URF2**Other Names**

E3 ubiquitin-protein ligase UHRF2, 632-, Np95/ICBP90-like RING finger protein, Np95-like RING finger protein, Nuclear protein 97, Nuclear zinc finger protein Np97, RING finger protein 107, Ubiquitin-like PHD and RING finger domain-containing protein 2, Ubiquitin-like-containing PHD and RING finger domains protein 2, UHRF2, NIRF, RNF107

Target/Specificity

UHRF2 encodes a nuclear protein which is involved in cell-cycle regulation. The encoded protein is a ubiquitin-ligase capable of ubiquinating PCNP (PEST-containing nuclear protein), and together they may play a role in tumorigenesis. This gene encodes a nuclear protein which is involved in cell-cycle regulation. The encoded protein is a ubiquitin-ligase capable of ubiquinating PCNP (PEST-containing nuclear protein), and together they may play a role in tumorigenesis.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-UHRF2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

UHRF2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

UHRF2 antibody - middle region - Protein Information

Name UHRF2**Synonyms** NIRF, RNF107**Function**

E3 ubiquitin ligase that plays important roles in DNA methylation, histone modifications, cell cycle and DNA repair (PubMed:15178429, PubMed:29506131, PubMed:27743347, PubMed:23404503). Acts as a specific reader for 5-hydroxymethylcytosine (5hmC) and thereby recruits various substrates to these sites to ubiquitinate them (PubMed:27129234, PubMed:24813944). This activity also allows the maintenance of 5mC levels at specific genomic loci and regulates neuron-related gene expression (By similarity). Participates in cell cycle regulation by ubiquitinating cyclins CCND1 and CCNE1 and thereby inducing G1 arrest (PubMed:15178429, PubMed:15361834, PubMed:21952639). Ubiquitinates also PCNP leading to its degradation by the proteasome (PubMed:14741369, PubMed:12176013). Plays an active role in DNA damage repair by ubiquitinating p21/CDKN1A leading to its proteasomal degradation (PubMed:29923055). Promotes also DNA repair by acting as an interstrand cross-links (ICLs) sensor. Mechanistically, cooperates with UHRF1 to ensure recruitment of FANCD2 to ICLs, leading to FANCD2 monoubiquitination and subsequent activation (PubMed:30335751). Contributes to UV-induced DNA damage response by physically interacting with ATR in response to irradiation, thereby promoting ATR activation (PubMed:33848395).

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00358, ECO:0000269|PubMed:12176013, ECO:0000269|PubMed:23404503, ECO:0000269|PubMed:27129234, ECO:0000269|PubMed:27743347, ECO:0000269|PubMed:29923055, ECO:0000269|PubMed:30335751}. Chromosome. Note=Enriched at genomic loci that are enriched for 5-hydroxymethylcytosine (5hmC)

UHRF2 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UHRF2 antibody - middle region - Images



UHRF2 antibody - middle region (AI10140) in Human Liver cells using Western Blot
WB Suggested Anti-UHRF2 Antibody Titration: 0.2-1 $\mu\text{g/ml}$
ELISA Titer: 1:62500
Positive Control: Human Liver

UHRF2 antibody - middle region - Background

This is a rabbit polyclonal antibody against UHRF2. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).