

TSC22D4 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10169**Specification****TSC22D4 antibody - N-terminal region - Product Information**

Application	WB, IHC
Primary Accession	O9Y3Q8
Other Accession	NM_030935 , NP_112197
Reactivity	Human, Rat, Dog
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

TSC22D4 antibody - N-terminal region - Additional Information**Gene ID** 81628**Alias Symbol** **THG1, THG-1, TILZ2****Other Names**

TSC22 domain family protein 4, TSC22-related-inducible leucine zipper protein 2, Tsc-22-like protein THG-1, TSC22D4, THG1, TILZ2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-TSC22D4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

TSC22D4 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TSC22D4 antibody - N-terminal region - Protein Information**Name** TSC22D4 ([HGNC:21696](#))**Function**

Binds DNA and acts as a transcriptional repressor (PubMed:[10488076](http://www.uniprot.org/citations/10488076)). Involved in the regulation of systematic glucose homeostasis and insulin sensitivity, via transcriptional repression of downstream insulin signaling targets such as OBP2A/LCN13 (By similarity). Acts as a negative regulator of lipogenic gene expression in hepatocytes and thereby mediates the control of very low-density lipoprotein release (PubMed:[23307490](http://www.uniprot.org/citations/23307490)). May play a role in neurite elongation and survival (By similarity).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9EQN3}. Cytoplasm {ECO:0000250|UniProtKB:Q9EQN3}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q9EQN3}. Synapse {ECO:0000250|UniProtKB:Q9EQN3} Note=Localizes away from the nucleus to neurite processes and synaptic termini as cerebellar granular neurons differentiate (By similarity) Accumulates in the cytoplasm of differentiated Purkinje cells (By similarity). Localized to both the cytoplasm and nucleus in immature cerebellar granular neurons and atrophic Purkinje cells (By similarity). {ECO:0000250|UniProtKB:Q9EQN3}

Tissue Location

Expressed in the liver.

TSC22D4 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

