

**TANK Antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI10174****Specification**

---

**TANK Antibody - N-terminal region - Product Information**

|                   |                                                 |
|-------------------|-------------------------------------------------|
| Application       | WB                                              |
| Primary Accession | <a href="#">E7EVA2</a>                          |
| Reactivity        | Human, Mouse, Rat, Pig, Horse, Bovine, Dog      |
| Predicted         | Human, Mouse, Rat, Pig, Bovine, Guinea Pig, Dog |
| Host              | Rabbit                                          |
| Clonality         | Polyclonal                                      |
| Calculated MW     | 26kDa KDa                                       |

**TANK Antibody - N-terminal region - Additional Information****Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-TANK antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


**Precautions**

TANK Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**TANK Antibody - N-terminal region - Protein Information****TANK Antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**TANK Antibody - N-terminal region - Images**



**Host:**Rabbit  
**Target Name:**TANK  
**Sample Tissue:** Lung Tumor lysates  
**Antibody Dilution:**1.μg/ml