10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Bcl2l2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10180

Specification

Bcl2l2 Antibody - C-terminal region - Product Information

Application	**WB**
Primary Accession	P70345
Other Accession	NM_007537, NP_031563
Reactivity	**Human, Mouse, Rat, Pig, Horse, Bovine, Dog**
Predicted	**Rat, Pig, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**21kDa KDa**

Bcl2l2 Antibody - C-terminal region - Additional Information

Gene ID 12050

Alias Symbol	**AW048834, Bcl-w, Gtrgal2, Gtrosa41, bclw, c98**

Other Names
Bcl-2-like protein 2, Bcl2-L-2, Apoptosis regulator Bcl-W, c98, Bcl2l2, Bclw, Kiaa0271

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-Bcl2l2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
Bcl2l2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Bcl2l2 Antibody - C-terminal region - Protein Information

Name Bcl2l2

Synonyms Bclw, Kiaa0271

Function
Promotes cell survival. Blocks dexamethasone-induced apoptosis. Mediates survival of postmitotic Sertoli cells by suppressing death-promoting activity of BAX.

Cellular Location
Mitochondrion membrane; Peripheral membrane protein. Note=Loosely associated with the

mitochondrial membrane in healthy cells. During apoptosis, tightly bound to the membrane (By similarity).


Tissue Location
Expressed in almost all myeloid cell lines and in a wide range of tissues, with highest levels in brain, colon, and salivary gland

Bcl2l2 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Bcl2l2 Antibody - C-terminal region - Images



WB Suggested Anti-Bcl2l2 Antibody
Titration: 1. µg/ml
Positive Control: Mouse Testis