

Fkhl18 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10270**Specification****Fkhl18 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	Q61574
Other Accession	NM_010226 , NP_034356
Reactivity	Human, Mouse, Rat, Horse, Bovine
Predicted	Human, Mouse, Rat, Zebrafish, Pig, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35kDa KDa

Fkhl18 antibody - N-terminal region - Additional Information**Gene ID** 14239**Alias Symbol** FREAC10, Fkh3, Fkhl18**Other Names**

Forkhead box protein S1, Forkhead-like 18 protein, Forkhead-related transcription factor 10, FREAC-10, Transcription factor FKH-3, Foxs1, Fkh3, Fkhl18, Freac10

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Fkhl18 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Fkhl18 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Fkhl18 antibody - N-terminal region - Protein Information**Name** Foxs1**Synonyms** Fkh3, Fkhl18, Freac10**Function**

Transcriptional repressor that suppresses transcription from the FASLG, FOXO3 and FOXO4 promoters. May have a role in the organization of the testicular vasculature.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00089, ECO:0000269|PubMed:18288644}

Fkhl18 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

