

**Foxo1 Antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI10271****Specification**

---

**Foxo1 Antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">G3V7R4</a>
Other Accession	<a href="#">NM_001191846</a> , <a href="#">NP_001178775</a>
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Bovine, Dog
Predicted	Human, Mouse, Rat, Zebrafish, Pig, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	71kDa KDa

**Foxo1 Antibody - N-terminal region - Additional Information****Gene ID** 84482**Alias Symbol** **Fkhr, Foxo1a****Other Names**

Forkhead box protein O1, Forkhead box protein O1A, Forkhead in rhabdomyosarcoma, Foxo1, Foxo1a

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-Foxo1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

Foxo1 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**Foxo1 Antibody - N-terminal region - Protein Information****Name** Foxo1**Synonyms** Foxo1a**Function**

Transcription factor that is the main target of insulin signaling and regulates metabolic homeostasis in response to oxidative stress (By similarity). Binds to the insulin response element (IRE) with consensus sequence 5'-TT[G/A]TTTGG-3' and the related Daf-16 family binding element (DBE) with consensus sequence 5'-TT[G/A]TTTAC-3'. Activity suppressed by insulin (By similarity).

Main regulator of redox balance and osteoblast numbers and controls bone mass (By similarity). Orchestrates the endocrine function of the skeleton in regulating glucose metabolism (By similarity). Also acts as a key regulator of chondrogenic commitment of skeletal progenitor cells in response to lipid availability: when lipids levels are low, translocates to the nucleus and promotes expression of SOX9, which induces chondrogenic commitment and suppresses fatty acid oxidation (By similarity). Acts synergistically with ATF4 to suppress osteocalcin/BGLAP activity, increasing glucose levels and triggering glucose intolerance and insulin insensitivity (By similarity). Also suppresses the transcriptional activity of RUNX2, an upstream activator of osteocalcin/BGLAP (By similarity). Acts as an inhibitor of glucose sensing in pancreatic beta cells by acting as a transcription repressor and suppressing expression of PDX1 (By similarity). In hepatocytes, promotes gluconeogenesis by acting together with PPARGC1A and CEBPA to activate the expression of genes such as IGFBP1, G6PC1 and PCK1 (By similarity). Also promotes gluconeogenesis by directly promoting expression of PPARGC1A and G6PC1 (By similarity). Important regulator of cell death acting downstream of CDK1, PKB/AKT1 and STK4/MST1 (By similarity). Promotes neural cell death (By similarity). Mediates insulin action on adipose tissue (By similarity). Regulates the expression of adipogenic genes such as PPARG during preadipocyte differentiation and, adipocyte size and adipose tissue-specific gene expression in response to excessive calorie intake (By similarity). Regulates the transcriptional activity of GADD45A and repair of nitric oxide-damaged DNA in beta-cells (By similarity). Required for the autophagic cell death induction in response to starvation or oxidative stress in a transcription-independent manner (By similarity). Mediates the function of MLIP in cardiomyocytes hypertrophy and cardiac remodeling (PubMed:<a href="http://www.uniprot.org/citations/26436652" target="\_blank">26436652</a>). Regulates endothelial cell (EC) viability and apoptosis in a PPIA/CYPA-dependent manner via transcription of CCL2 and BCL2L11 which are involved in EC chemotaxis and apoptosis (By similarity).


#### Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q9R1E0}. Nucleus {ECO:0000250|UniProtKB:Q9R1E0}. Note=Shuttles between the cytoplasm and nucleus (By similarity). Largely nuclear in unstimulated cells (By similarity). In osteoblasts, colocalizes with ATF4 and RUNX2 in the nucleus. Serum deprivation increases localization to the nucleus, leading to activate expression of SOX9 and subsequent chondrogenesis (By similarity). Insulin-induced phosphorylation at Ser-253 by PKB/AKT1 leads, via stimulation of Thr-24 phosphorylation, to binding of 14-3-3 proteins and nuclear export to the cytoplasm where it is degraded by the ubiquitin-proteasomal pathway (By similarity). Phosphorylation at Ser-249 by CDK1 disrupts binding of 14-3-3 proteins and promotes nuclear accumulation (By similarity). Phosphorylation by NLK results in nuclear export (By similarity). Translocates to the nucleus upon oxidative stress-induced phosphorylation at Ser-212 by STK4/MST1 (By similarity). SGK1-mediated phosphorylation also results in nuclear translocation. Retained in the nucleus under stress stimuli including oxidative stress, nutrient deprivation or nitric oxide. Methylated form is nuclear (By similarity). PPIA/CYPA stimulates its nuclear accumulation (By similarity). Deacetylation by SIRT6, promotes its translocation into the cytoplasm (By similarity) {ECO:0000250|UniProtKB:Q12778, ECO:0000250|UniProtKB:Q9R1E0}

#### Foxo1 Antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Foxo1 Antibody - N-terminal region - Images****WB Suggested Anti-Foxo1 Antibody****Titration: 1. µg/ml**

Positive Control: Rat Spleen