

Skil antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10351**Specification**

Skil antibody - C-terminal region - Product Information

Application	WB
Primary Accession	D3ZWL1
Other Accession	XM_001057072 , XP_001057072
Reactivity	Human, Mouse, Rat, Pig, Horse, Yeast, Bovine, Dog
Predicted	Human, Mouse, Rat, Pig, Chicken, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	74kDa kDa

Skil antibody - C-terminal region - Additional Information

Alias Symbol **Skir, Sno**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Skil antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Skil antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Skil antibody - C-terminal region - Protein Information**Skil antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Skil antibody - C-terminal region - Images

**WB Suggested Anti-Skil Antibody****Titration: 1. µg/ml**

Positive Control: Rat Liver