

PATZ1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10374**Specification****PATZ1 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	Q9HBE1
Other Accession	NM_014323 , NP_055138
Reactivity	Human, Mouse, Rat, Horse, Bovine, Dog
Predicted	Human, Mouse, Pig, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	74kDa KDa

PATZ1 antibody - N-terminal region - Additional Information**Gene ID** 23598

Alias Symbol **MAZR, PATZ, RIAZ, ZBTB19, ZNF278, ZSG, dj400N23**

Other Names

POZ-, AT hook-, and zinc finger-containing protein 1, BTB/POZ domain zinc finger transcription factor, Protein kinase A RI subunit alpha-associated protein, Zinc finger and BTB domain-containing protein 19, Zinc finger protein 278, Zinc finger sarcoma gene protein, PATZ1, PATZ, RIAZ, ZBTB19, ZNF278, ZSG

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PATZ1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

PATZ1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PATZ1 antibody - N-terminal region - Protein Information

Name PATZ1

Synonyms PATZ, RIAZ, ZBTB19, ZNF278, ZSG

Function

Transcriptional regulator that plays a role in many biological processes such as embryogenesis, senescence, T-cell development or neurogenesis (PubMed:10713105, PubMed:<a

[25755280](http://www.uniprot.org/citations/25755280), PubMed: [31875552](http://www.uniprot.org/citations/31875552)). Interacts with the TP53 protein to control genes that are important in proliferation and in the DNA-damage response. Mechanistically, the interaction inhibits the DNA binding and transcriptional activity of TP53/p53 (PubMed: [25755280](http://www.uniprot.org/citations/25755280)). Part of the transcriptional network modulating regulatory T-cell development and controls the generation of the regulatory T-cell pool under homeostatic conditions (PubMed: [31875552](http://www.uniprot.org/citations/31875552)).

Cellular Location

Nucleus.

Tissue Location

Ubiquitous.

PATZ1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

