

**Atf2 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI10613****Specification****Atf2 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">Q00969</a>
Other Accession	<a href="#">NM_031018</a> , <a href="#">NP_112280</a>
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Pig, Chicken, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 52kDa KDa

**Atf2 antibody - N-terminal region - Additional Information****Gene ID** 81647**Other Names**

Cyclic AMP-dependent transcription factor ATF-2, cAMP-dependent transcription factor ATF-2, 2.3.1.48, Activating transcription factor 2, cAMP response element-binding protein CRE-BP1, Atf2

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-Atf2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

Atf2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**Atf2 antibody - N-terminal region - Protein Information****Name** Atf2**Function**

Transcriptional activator which regulates the transcription of various genes, including those involved in anti-apoptosis, cell growth, and DNA damage response. Dependent on its binding partner, binds to CRE (cAMP response element) consensus sequences (5'-TGACGTCA- 3') or to AP-1 (activator protein 1) consensus sequences (5'-TGACTCA- 3'). In the nucleus, contributes to global transcription and the DNA damage response, in addition to specific transcriptional activities that are related to cell development, proliferation and death. In the cytoplasm, interacts with and perturbs HK1- and VDAC1-containing complexes at the mitochondrial outer membrane, thereby impairing mitochondrial membrane potential, inducing mitochondrial leakage and promoting cell

death. The phosphorylated form (mediated by ATM) plays a role in the DNA damage response and is involved in the ionizing radiation (IR)-induced S phase checkpoint control and in the recruitment of the MRN complex into the IR-induced foci (IRIF). Exhibits histone acetyltransferase (HAT) activity which specifically acetylates histones H2B and H4 in vitro. In concert with CUL3 and RBX1, promotes the degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM. Can elicit oncogenic or tumor suppressor activities depending on the tissue or cell type (By similarity).

#### **Cellular Location**

Nucleus. Cytoplasm. Mitochondrion outer membrane. Note=Shuttles between the cytoplasm and the nucleus and heterodimerization with JUN is essential for the nuclear localization. Localization to the cytoplasm is observed under conditions of cellular stress and in disease states. Localizes at the mitochondrial outer membrane in response to genotoxic stress Phosphorylation at Thr-34 is required for its nuclear localization and negatively regulates its mitochondrial localization. Colocalizes with the MRN complex in the IR-induced foci (IRIF) (By similarity)

#### **Atf2 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

