

ZSCAN20 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10662**Specification**

ZSCAN20 antibody - middle region - Product Information

Application	WB
Primary Accession	P17040
Other Accession	NM_145238 , NP_660281
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted Host	Rat, Horse, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 117kDa KDa

ZSCAN20 antibody - middle region - Additional Information**Gene ID 7579****Alias Symbol** **KOX29, ZFP-31, ZNF31, ZNF360****Other Names**

Zinc finger and SCAN domain-containing protein 20, Zinc finger protein 31, Zinc finger protein 360, Zinc finger protein KOX29, ZSCAN20, KOX29, ZNF31, ZNF360

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZSCAN20 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZSCAN20 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

ZSCAN20 antibody - middle region - Protein Information**Name** ZSCAN20**Synonyms** KOX29, ZNF31, ZNF360**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187}.

ZSCAN20 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZSCAN20 antibody - middle region - Images



WB Suggested Anti-ZSCAN20 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Hela cell lysate

ZSCAN20 antibody - middle region - References

Jordanova, A., (2006) Nat. Genet. 38 (2), 197-202 Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.