

TBC1D10B antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10680**Specification**

TBC1D10B antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q9H8Z2
Other Accession	NM_015527 , NP_056342
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Yeast, Bovine, Dog
Predicted Host	Horse, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 61kDa KDa

TBC1D10B antibody - C-terminal region - Additional Information

Alias Symbol **EPI64B, FP2461**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-TBC1D10B antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


TBC1D10B antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TBC1D10B antibody - C-terminal region - Protein Information**TBC1D10B antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TBC1D10B antibody - C-terminal region - Images



WB Suggested Anti-TBC1D10B Antibody Titration: 1.25µg/ml

ELISA Titer: 1:12500

Positive Control: HepG2 cell lysate

TBC1D10B is supported by BioGPS gene expression data to be expressed in HepG2

TBC1D10B antibody - C-terminal region - References

Ota,T., et al., (2004) Nat. Genet. 36 (1), 40-45
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.