10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ZNF396 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10746

Specification

ZNF396 antibody - middle region - Product Information

Application	**WB**
Primary Accession	Q96N95
Other Accession	NM_145756, NP_665699
Reactivity	**Human, Rabbit**
Predicted	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**38kDa KDa**

ZNF396 antibody - middle region - Additional Information

Gene ID 252884

Alias Symbol	**FLJ31213, ZSCAN14**

Other Names
Zinc finger protein 396, Zinc finger and SCAN domain-containing protein 14, ZNF396, ZSCAN14

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-ZNF396 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
ZNF396 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF396 antibody - middle region - Protein Information

Name ZNF396

Synonyms ZSCAN14

Function
Isoform 1 and isoform 2 act as DNA-dependent transcriptional repressors.

Cellular Location
[Isoform 1]: Nucleus.

Tissue Location
Expressed strongly in liver, moderately in skeletal muscle and weakly in kidney, pancreas, spleen


and prostate

ZNF396 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ZNF396 antibody - middle region - Images



WB Suggested Anti-ZNF396 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:62500
Positive Control: Jurkat cell lysate

ZNF396 antibody - middle region - References

Wu,Y., Gene 310, 193-201 (2003)Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.