

DIDO1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10750**Specification**

DIDO1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q9BTC0-2
Other Accession	NM_080797 , NP_542987
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Yeast, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Chicken, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59kDa KDa

DIDO1 antibody - N-terminal region - Additional Information

Alias Symbol	BYE1, DIO1, DATF1, DIDO2, DIDO3, DIO-1, DATF-1, C20orf158, dj885L7.8
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DIDO1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

DIDO1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DIDO1 antibody - N-terminal region - Protein Information**DIDO1 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DIDO1 antibody - N-terminal region - Images

WB Suggested Anti-DIDO1 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: Jurkat cell lysate

DIDO1 is strongly supported by BioGPS gene expression data to be expressed in Human Jurkat cells