

CACNB4 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10766**Specification**

CACNB4 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O00305-2
Other Accession	NM_001005747 , NP_001005747
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55kDa KDa

CACNB4 antibody - C-terminal region - Additional Information

Alias Symbol	CAB4, CACNLB4, EA5, EJM, EIG9, EJM4, EJM6
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CACNB4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

CACNB4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CACNB4 antibody - C-terminal region - Protein Information**CACNB4 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CACNB4 antibody - C-terminal region - Images

WB Suggested Anti-CACNB4 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:12500

Positive Control: Jurkat cell lysate

CACNB4 antibody - C-terminal region - References

Suzuki,Y., et al., (2004) Genome Res. 14 (9), 1711-1718
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.