

GRIK2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10821**Specification**

GRIK2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O96KS6
Other Accession	NM_175768 , NP_786944
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	98kDa KDa

GRIK2 antibody - C-terminal region - Additional Information

Alias Symbol	EAA4, GLR6, MRT6, GLUK6, GLUR6, GluK2
Format	
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.	

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-GRIK2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


GRIK2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

GRIK2 antibody - C-terminal region - Protein Information**GRIK2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GRIK2 antibody - C-terminal region - Images



WB Suggested Anti-GRIK2 Antibody Titration: 0.125µg/ml

ELISA Titer: 1:62500

Positive Control: Human brain

GRIK2 antibody - C-terminal region - References

Bah,J., et al., (2004) Neuroreport 15 (12), 1987-1991
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.