

SCAND1 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10910**Specification**

SCAND1 antibody - middle region - Product Information

Application	WB
Primary Accession	P57086
Other Accession	NM_016558 , NP_057642
Reactivity	Human, Pig
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19kDa KDa

SCAND1 antibody - middle region - Additional Information**Gene ID** 51282**Alias Symbol** **RAZ1, SDP1****Other Names**

SCAN domain-containing protein 1, SCAND1, SDP1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SCAND1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SCAND1 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

SCAND1 antibody - middle region - Protein Information**Name** SCAND1**Synonyms** SDP1**Function**

May regulate transcriptional activity.

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187}.

SCAND1 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SCAND1 antibody - middle region - Images



WB Suggested Anti-SCAND1 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

SCAND1 is supported by BioGPS gene expression data to be expressed in HepG2

SCAND1 antibody - middle region - References

Carneiro, F.R., et al., (2006) Biochem. Biophys. Res. Commun. 343 (1), 260-268
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.