10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ZNF167 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10916

Specification

ZNF167 antibody - N-terminal region - Product Information

Application	**IHC, WB**
Primary Accession	Q9P0L1
Other Accession	NM_018651, NP_061121
Reactivity	**Human, Rat**
Predicted	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**85kDa KDa**

ZNF167 antibody - N-terminal region - Additional Information

Gene ID 55888

Alias Symbol **ZFP, ZNF64, ZNF448, ZKSCAN7, ZNF167**

Other Names
Zinc finger protein with KRAB and SCAN domains 7, Zinc finger protein 167, Zinc finger protein 448, Zinc finger protein 64, ZKSCAN7, ZNF167, ZNF448, ZNF64

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 100 ul of distilled water. Final anti-ZNF167 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
ZNF167 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF167 antibody - N-terminal region - Protein Information

Name ZKSCAN7

Synonyms ZNF167, ZNF448, ZNF64

Function
May be involved in transcriptional regulation.

Cellular Location
Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187}.


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ZNF167 antibody - N-terminal region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ZNF167 antibody - N-terminal region - Images



Human Intestine



WB Suggested Anti-ZNF167 Antibody Titration: 1.25µg/ml
ELISA Titer: 1:312500
Positive Control: Jurkat cell lysate

ZNF167 antibody - N-terminal region - References

Calabro,V., et al., (1995) Hum. Genet. 95 (1), 18-21Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.