

SCAND2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10922**Specification**

SCAND2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O9GZW5
Other Accession	NM_022050 , NP_071333
Reactivity	Human
Predicted	Human, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa KDa

SCAND2 antibody - N-terminal region - Additional Information

Alias Symbol **SCAND2,**
Other Names
Putative SCAN domain-containing protein SCAND2P, SCAN domain-containing protein 2
pseudogene, SCAND2P, SCAND2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SCAND2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SCAND2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SCAND2 antibody - N-terminal region - Protein Information

Name SCAND2P

Synonyms SCAND2

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187}.

SCAND2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SCAND2 antibody - N-terminal region - Images



WB Suggested Anti-SCAND2 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:1562500

Positive Control: MCF7 cell lysate