

GIOT-1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10954**Specification**

GIOT-1 antibody - N-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q8TAF7-2
Other Accession	NM_153257 , AAH28631
Reactivity	Human, Horse, Dog
Predicted	Human, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48kDa KDa

GIOT-1 antibody - N-terminal region - Additional Information

Alias Symbol **GIOT1, Giot-1**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-GIOT-1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


GIOT-1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

GIOT-1 antibody - N-terminal region - Protein Information**GIOT-1 antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GIOT-1 antibody - N-terminal region - Images



Human Intestine



WB Suggested Anti-GIOT-1 Antibody Titration: 1.25µg/ml

ELISA Titer: 1:312500

Positive Control: Human Thymus