

HMG1L10 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10955**Specification**

HMG1L10 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q9UGV6
Other Accession	CAB62951
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23kDa KDa

HMG1L10 antibody - C-terminal region - Additional Information

Alias Symbol **bK445C9.3**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-HMG1L10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


HMG1L10 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HMG1L10 antibody - C-terminal region - Protein Information**HMG1L10 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HMG1L10 antibody - C-terminal region - Images



WB Suggested Anti-HMG1L10 Antibody Titration: 1.25µg/ml
ELISA Titer: 1:62500
Positive Control: HepG2 cell lysate