

DDX19A antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10985**Specification**

DDX19A antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q9NUU7
Other Accession	NM_018332 , NP_060802
Reactivity	Human, Mouse, Rat, Horse, Dog
Predicted	Rat, Rabbit, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53kDa KDa

DDX19A antibody - N-terminal region - Additional Information**Gene ID** 55308**Alias Symbol** **DDX19L, DDX19-DDX19L****Other Names**

ATP-dependent RNA helicase DDX19A, 3.6.4.13, DDX19-like protein, DEAD box protein 19A, DDX19A, DDX19L

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-DDX19A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

DDX19A antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DDX19A antibody - N-terminal region - Protein Information**Name** DDX19A**Synonyms** DDX19L**Function**

ATP-dependent RNA helicase involved in mRNA export from the nucleus. Rather than unwinding RNA duplexes, DDX19 functions as a remodeler of ribonucleoprotein particles, whereby proteins bound to nuclear mRNA are dissociated and replaced by cytoplasmic mRNA binding proteins.

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q9UMR2}. Nucleus, nucleoplasm


{ECO:0000250|UniProtKB:Q9UMR2}. Note=Associates with the nuclear pore complex cytoplasmic fibrils {ECO:0000250|UniProtKB:Q9UMR2}

DDX19A antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DDX19A antibody - N-terminal region - Images



WB Suggested Anti-DDX19A Antibody Titration: 2.5µg/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

DDX19A is strongly supported by BioGPS gene expression data to be expressed in Human HepG2 cells