

HSF1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11059**Specification**

HSF1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P38532-2
Other Accession	NM_008296 , NP_032322
Reactivity	Human, Mouse, Rat, Rabbit, Goat, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Pig, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 55kDa KDa

HSF1 antibody - C-terminal region - Additional Information

Alias Symbol **AA960185**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-HSF1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


HSF1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HSF1 antibody - C-terminal region - Protein Information**HSF1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HSF1 antibody - C-terminal region - Images



WB Suggested Anti-HSF1 Antibody Titration: 5.0µg/ml

ELISA Titer: 1:62500

Positive Control: NIH/3T3 cell lysate

HSF1 antibody - C-terminal region - References

Shamovsky, I., et al., (2006) Nature 440 (7083), 556-560
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.