

Hsf2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11060**Specification**

Hsf2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8BWK6
Other Accession	NM_008297 , NP_032323
Reactivity	Human, Mouse, Rat, Rabbit
Predicted	Human, Mouse, Rat, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	58kDa KDa

Hsf2 antibody - C-terminal region - Additional Information

Alias Symbol **AI661205**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Hsf2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Hsf2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Hsf2 antibody - C-terminal region - Protein Information**Hsf2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Hsf2 antibody - C-terminal region - Images



90 kDa__
65 kDa__
40 kDa__
31 kDa__
22 kDa__

WB Suggested Anti-Hsf2 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:312500

Positive Control: Mouse Uterus