

Rel antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11079**Specification**

Rel antibody - middle region - Product Information

Application	WB
Primary Accession	A4QPD3
Other Accession	NM_009044 , NP_033070
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Pig, Horse, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 65kDa KDa

Rel antibody - middle region - Additional Information

Alias Symbol **c-Rel**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Rel antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Rel antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Rel antibody - middle region - Protein Information**Rel antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Rel antibody - middle region - Images



WB Suggested Anti-Rel Antibody Titration: 0.2-1 μ g/ml
Positive Control: Mouse Lung