

Hoxa1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11116**Specification**

Hoxa1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	B9EHK7
Other Accession	NM_010449 , NP_034579
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Predicted	Mouse, Rabbit, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36 KDa

Hoxa1 antibody - C-terminal region - Additional Information

Alias Symbol **ERA1, Hox-1.6**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Hoxa1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Hoxa1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Hoxa1 antibody - C-terminal region - Protein Information**Hoxa1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Hoxa1 antibody - C-terminal region - Images



WB Suggested Anti-Hoxa1 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Mouse Small Intestine