

Dmrt1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11153**Specification**

Dmrt1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q6Y951
Other Accession	NM_015826 , NP_056641
Reactivity	Mouse, Rat
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39kDa KDa

Dmrt1 antibody - C-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Dmrt1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


Dmrt1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Dmrt1 antibody - C-terminal region - Protein Information**Dmrt1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Dmrt1 antibody - C-terminal region - Images



WB Suggested Anti-Dmrt1 Antibody Titration: 0.2-1 μ g/ml
Positive Control: NIH/3T3 cell lysate