

PUS1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11168**Specification**

PUS1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q791J1
Other Accession	NM_001025562 , NP_001020733
Reactivity	Human, Mouse, Rat
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43kDa KDa

PUS1 antibody - C-terminal region - Additional Information

Alias Symbol **MPUS1, mPus1p, A730013B20Rik**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PUS1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


PUS1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PUS1 antibody - C-terminal region - Protein Information**PUS1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PUS1 antibody - C-terminal region - Images



WB Suggested Anti-PUS1 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:312500

Positive Control: SP2/0 cell lysate

PUS1 antibody - C-terminal region - References

Zhao,X., et al., (2004) Mol. Cell 15 (4), 549-558
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.