

BRF1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11204**Specification**

BRF1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q6PED6
Other Accession	NM_028193 , AAH58112
Reactivity	Mouse, Rat, Yeast, Bovine, Dog
Predicted	Mouse, Rat, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22kDa KDa

BRF1 antibody - C-terminal region - Additional Information

Alias Symbol	GTF3B, TAF3C, TAFIII90, TFIIB90, mTFIIB90, 2510002F24Rik
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-BRF1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


BRF1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

BRF1 antibody - C-terminal region - Protein Information**BRF1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BRF1 antibody - C-terminal region - Images



WB Suggested Anti-BRF1 Antibody Titration: 1.25µg/ml

ELISA Titer: 1:62500

Positive Control: NIH/3T3 cell lysate

BRF1 antibody - C-terminal region - References

Strausberg, R.L., et al., (2002) Proc. Natl. Acad. Sci. U.S.A. 99 (26), 16899-16903
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.