

**Tfdp2 antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI11262****Specification**

---

**Tfdp2 antibody - C-terminal region - Product Information**

|                   |                                                       |
|-------------------|-------------------------------------------------------|
| Application       | WB                                                    |
| Primary Accession | <a href="#">Q8BHD2</a>                                |
| Other Accession   | <a href="#">NM_178667</a> , <a href="#">NP_848782</a> |
| Reactivity        | Human, Mouse, Rat, Pig, Horse, Bovine, Dog            |
| Predicted Host    | Human, Mouse, Rat, Horse, Bovine, Dog                 |
| Clonality         | Rabbit                                                |
| Calculated MW     | Polyclonal<br>43kDa KDa                               |

**Tfdp2 antibody - C-terminal region - Additional Information**

Alias Symbol **A330080J22Rik, DP-3, DP3, 1110029I05Rik**

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-Tfdp2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

Tfdp2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


**Tfdp2 antibody - C-terminal region - Protein Information****Tfdp2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Tfdp2 antibody - C-terminal region - Images**

---



WB Suggested Anti-Tfdp2 Antibody Titration: 0.2-1  $\mu$ g/ml  
Positive Control: Mouse Uterus