

Foxi2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11271**Specification**

Foxi2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	A2RTG9
Other Accession	NM_183193 , NP_899016
Reactivity	Mouse
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa KDa

Foxi2 antibody - C-terminal region - Additional Information

Alias Symbol **B130055A05Rik**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Foxi2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

Foxi2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Foxi2 antibody - C-terminal region - Protein Information**Foxi2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Foxi2 antibody - C-terminal region - Images



90 kDa
65 kDa
40 kDa
31 kDa
22 kDa

WB Suggested Anti-Foxi2 Antibody Titration: 0.2-1 μ g/ml
Positive Control: NIH/3T3 cell lysate