

POU2F1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11280**Specification**

POU2F1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P25425-10
Other Accession	NM_198932 , NP_945150
Reactivity	Mouse
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57kDa KDa

POU2F1 antibody - C-terminal region - Additional Information

Alias Symbol	Oct1, Otf1, NF-A1, Oct-1, Otf-1, 2810482H01Rik
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-POU2F1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


POU2F1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

POU2F1 antibody - C-terminal region - Protein Information**POU2F1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

POU2F1 antibody - C-terminal region - Images



WB Suggested Anti-POU2F1 Antibody Titration: 1.25µg/ml
ELISA Titer: 1:62500
Positive Control: SP2/0 cell lysate

POU2F1 antibody - C-terminal region - References

Tantin,D., et al., (2005) Cancer Res. 65 (23), 10750-10758
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.
Publications: Gupta, S. et al. Human organic cation transporter 1 is expressed in lymphoma cells and increases susceptibility to irinotecan and paclitaxel. J. Pharmacol. Exp. Ther. 341, 16-23 (2012). WB, Mouse22202118