

ZHX2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11282**Specification**

ZHX2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8°C0C0
Other Accession	NM_199449 , NP_955520
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	92kDa KDa

ZHX2 antibody - C-terminal region - Additional Information

Alias Symbol **Raf, Afr1, Afr-1, mKIAA0854**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ZHX2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


ZHX2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZHX2 antibody - C-terminal region - Protein Information**ZHX2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZHX2 antibody - C-terminal region - Images



WB Suggested Anti-ZHX2 Antibody Titration: 2.5µg/ml

ELISA Titer: 1:62500

Positive Control: SP2/0 cell lysate

ZHX2 antibody - C-terminal region - References

Perincheri, S., et al., (2005) Proc. Natl. Acad. Sci. U.S.A. 102 (2), 396-401
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.