10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Naca-rs antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11291

Specification

Naca-rs antibody - middle region - Product Information

Application	**WB**
Other Accession	XM_354689, XP_354689
Reactivity	**Mouse**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**31kDa KDa**

Naca-rs antibody - middle region - Additional Information

Alias Symbol	**Gm1878**

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-Naca-rs antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
Naca-rs antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Naca-rs antibody - middle region - Protein Information


Naca-rs antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Naca-rs antibody - middle region - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



WB Suggested Anti-Naca-rs Antibody Titration: 0.2-1 µg/ml
Positive Control: Mouse Spleen