

Vgll2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11335**Specification**

Vgll2 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8BGW8
Other Accession	NM_153786 , NP_722481
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa kDa

Vgll2 Antibody - C-terminal region - Additional Information**Gene ID** 215031**Alias Symbol** **C130057C21Rik, VITO-1, Vito1****Other Names**

Transcription cofactor vestigial-like protein 2, Vgl-2, Protein VITO1, Vgll2, Vgl2, Vito1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Vgll2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Vgll2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Vgll2 Antibody - C-terminal region - Protein Information**Name** Vgll2**Synonyms** Vgl2, Vito1**Function**

May act as a specific coactivator for the mammalian TEFs. May play a role in the development of skeletal muscles.

Cellular Location

Nucleus.


Tissue Location

Skeletal muscle specific.

Vgll2 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Vgll2 Antibody - C-terminal region - Images

Host: Rabbit

Target Name: Vgll2

Sample Tissue: Mouse Thymus lysates

Antibody Dilution: 1.0µg/ml