

ENO1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11352**Specification**

ENO1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q53FT9
Other Accession	NM_001428 , NP_001419
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Sheep
Predicted	Rabbit, Zebrafish
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47kDa KDa

ENO1 antibody - C-terminal region - Additional Information

Alias Symbol **NNE, PPH, MPB1, ENO1L1**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ENO1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


ENO1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ENO1 antibody - C-terminal region - Protein Information**ENO1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ENO1 antibody - C-terminal region - Images



WB Suggested Anti-ENO1 Antibody Titration: 1.25µg/ml

ELISA Titer: 1:62500

Positive Control: Jurkat cell lysate

ENO1 is strongly supported by BioGPS gene expression data to be expressed in Human Jurkat cells

ENO1 antibody - C-terminal region - References

Ghosh,A.K., et al., (2005) J. Biol. Chem. 280 (14), 14325-14330
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.