

E2F3 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11359**Specification**

E2F3 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q68DT0
Other Accession	NM_001243076 , CAH18140
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rabbit, Pig, Chicken, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37kDa KDa

E2F3 antibody - C-terminal region - Additional Information

Alias Symbol **E2F-3**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-E2F3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


E2F3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

E2F3 antibody - C-terminal region - Protein Information**E2F3 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

E2F3 antibody - C-terminal region - Images



WB Suggested Anti-E2F3 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:62500
Positive Control: Human Thymus

E2F3 antibody - C-terminal region - References

Koehrer, K., et al., (2004) The German cDNA Consortium Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles. Publications: Naga Prasad, S. V et al. Unique microRNA profile in end-stage heart failure indicates alterations in specific cardiovascular signaling networks. J. Biol. Chem. 284, 27487-99 (2009). WB, IHC, Rat, Bovine, Pig, H, Guinea pig, Dog, Human, Mouse, Rabbit19641226 Melcher, R. et al. LOH and copy neutral LOH (cnLOH) act as alternative mechanism in sporadic colorectal cancers with chromosomal and microsatellite instability. Carcinogenesis 32, 636-42 (2011). WB, IHC, Rat, Bovine, Pig, H, Guinea pig, Dog, Human, Mouse, Rabbit21297112 Giangreco, A. A. et al. Tumor suppressor microRNAs, miR-100 and -125b, are regulated by 1,25-dihydroxyvitamin D in primary prostate cells and in patient tissue. Cancer Prev. Res. (Phila). 6, 483-94 (2013). WB, IHC, Rat, Bovine, Pig, H, Guinea pig, Dog, Human, Mouse, Rabbit23503652