

NR2C1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11387**Specification**

NR2C1 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q15625
Other Accession	NM_003297 , NP_003288
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67kDa KDa

NR2C1 antibody - C-terminal region - Additional Information

Alias Symbol **TR2, TR2-11**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NR2C1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


NR2C1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NR2C1 antibody - C-terminal region - Protein Information**NR2C1 antibody - C-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NR2C1 antibody - C-terminal region - Images



Human Lung



WB Suggested Anti-NR2C1 Antibody Titration: 0.1µg/ml

Positive Control: HepG2 cell lysate

NR2C1 is strongly supported by BioGPS gene expression data to be expressed in Human HepG2 cells

NR2C1 antibody - C-terminal region - References

Li, G., (2003) Biochem. Biophys. Res. Commun. 310 (2), 384-390
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.