

HCFC1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11436**Specification**

HCFC1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	P51610-3
Other Accession	NM_005334 , AAH63435
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	45kDa KDa

HCFC1 antibody - N-terminal region - Additional Information

Alias Symbol **CFF, HCF1, HFC1, VCAF, HCF-1**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-HCFC1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


HCFC1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HCFC1 antibody - N-terminal region - Protein Information**HCFC1 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HCFC1 antibody - N-terminal region - Images



WB Suggested Anti-HCFC1 Antibody Titration: 1.25µg/ml

Positive Control: HepG2 cell lysate

HCFC1 is strongly supported by BioGPS gene expression data to be expressed in Human HepG2 cells

HCFC1 antibody - N-terminal region - References

Director MGC Project, (2003) Direct Submission Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.