

TAF1C antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11442**Specification**

TAF1C antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q15572-2
Other Accession	NM_139353 , NP_647610
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85kDa KDa

TAF1C antibody - C-terminal region - Additional Information

Alias Symbol **SL1, TAFI95, TAFI110, MGC:39976**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-TAF1C antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


TAF1C antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TAF1C antibody - C-terminal region - Protein Information**TAF1C antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TAF1C antibody - C-terminal region - Images



WB Suggested Anti-TAF1C Antibody Titration: 2.5µg/ml

ELISA Titer: 1:12500

Positive Control: Jurkat cell lysate

TAF1C is strongly supported by BioGPS gene expression data to be expressed in Human Jurkat cells

TAF1C antibody - C-terminal region - References

Kimura, K., et al., (2006) Genome Res. 16 (1), 55-65
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.