

Pura antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11443**Specification**

Pura antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P42669
Other Accession	NM_008989 , NP_033015
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Sheep, Yeast, Bovine, Dog
Predicted Host	Mouse, Rat, Rabbit, Sheep, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 35kDa kDa

Pura antibody - C-terminal region - Additional Information**Gene ID** 19290**Alias Symbol** CAGER-1, Pur-alpha, ssCRE-BP**Other Names**

Transcriptional activator protein Pur-alpha, Purine-rich single-stranded DNA-binding protein alpha, Pura

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Pura antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Pura antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Pura antibody - C-terminal region - Protein Information**Name** Pura**Function**

This is a probable transcription activator that specifically binds the purine-rich single strand of the PUR element located upstream of the c-Myc gene. May play a role in the initiation of DNA replication and in recombination.

Cellular Location


Nucleus.

Pura antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Pura antibody - C-terminal region - Images



WB Suggested Anti-Pura Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Thymus