

JMJD2D antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11528**Specification**

JMJD2D antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q0VF39
Other Accession	NM_018039 , NP_060509
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59kDa KDa

JMJD2D antibody - C-terminal region - Additional Information

Alias Symbol **FLJ10251, JMJD2D**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-JMJD2D antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


JMJD2D antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

JMJD2D antibody - C-terminal region - Protein Information**JMJD2D antibody - C-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


JMJD2D antibody - C-terminal region - Images



Human Intestine



Human kidney



WB Suggested Anti-JMJD2D Antibody Titration: 0.2-1 μ g/ml
Positive Control: Human brain

JMJD2D antibody - C-terminal region - References

Katoh,M. (2004) Int. J. Oncol. 24 (6), 1623-1628
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.