

ZNF70 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11568**Specification**

ZNF70 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	O9UC06
Other Accession	NM_021916 , NP_068735
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted Host	Human, Horse, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 51kDa KDa

ZNF70 antibody - C-terminal region - Additional Information**Gene ID** 7621

Alias Symbol	Cos17
Other Names	
Zinc finger protein 70, Zinc finger protein N27C7-1, ZNF70	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ZNF70 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF70 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF70 antibody - C-terminal region - Protein Information**Name** ZNF70**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

ZNF70 antibody - C-terminal region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF70 antibody - C-terminal region - Images



Human Stomach



WB Suggested Anti-ZNF70 Antibody Titration: 5.0µg/ml

ELISA Titer: 1:312500

Positive Control: HepG2 cell lysate

ZNF70 antibody - C-terminal region - References

Collins, J.E., Genome Biol. 5 (10), R84 (2004) Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.