

NR1I2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11571**Specification**

NR1I2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9UNW4
Other Accession	NM_022002 , NP_071285
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Pig, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 54kDa KDa

NR1I2 antibody - C-terminal region - Additional Information

Alias Symbol	BXR, PAR, PRR, PXR, SAR, SXR, ONR1, PAR1, PAR2, PARq
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-NR1I2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


NR1I2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NR1I2 antibody - C-terminal region - Protein Information**NR1I2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NR1I2 antibody - C-terminal region - Images



WB Suggested Anti-NR1I2 Antibody Titration: 5.0µg/ml

ELISA Titer: 1:12500

Positive Control: Jurkat cell lysate

NR1I2 antibody - C-terminal region - References

Dring, M.M., (2006) Gastroenterology 130 (2), 341-348
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.