

Zkscan14 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11600**Specification**

Zkscan14 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9Z1D9
Other Accession	NM_023322 , NP_075811
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59kDa KDa

Zkscan14 antibody - C-terminal region - Additional Information**Gene ID** 67235**Alias Symbol** 2310046C23Rik, 2810437E14Rik, MGC117641, Zfp99, Znf394**Other Names**

Zinc finger protein 394, Zinc finger protein 94, Zfp-94, Zinc finger protein with KRAB and SCAN domains 14, Znf394, Zfp94, Zfp99, Zkscan14

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Zkscan14 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Zkscan14 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Zkscan14 antibody - C-terminal region - Protein Information**Name** Znf394**Synonyms** Zfp94, Zfp99, Zkscan14**Function**

May be involved in transcriptional regulation.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187}.


Tissue Location

Expressed at high level in testis.

Zkscan14 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Zkscan14 antibody - C-terminal region - Images

WB Suggested Anti-Zkscan14 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Brain