10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# ZNF485 antibody - N-terminal region
**Rabbit Polyclonal Antibody**
**Catalog # AI11635**

## Specification

### ZNF485 antibody - N-terminal region - Product Information

| | |
|---|---|
| Application | **WB** |
| Primary Accession | Q8NCK3 |
| Other Accession | NM_145312, NP_660355 |
| Reactivity | **Human, Mouse, Rat, Rabbit, Dog** |
| Predicted | **Human, Dog** |
| Host | **Rabbit** |
| Clonality | **Polyclonal** |
| Calculated MW | **46kDa KDa** |

### ZNF485 antibody - N-terminal region - Additional Information

**Gene ID** 220992

**Other Names**
Zinc finger protein 485, ZNF485

**Format**
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**
Add 50 ul of distilled water. Final anti-ZNF485 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**
ZNF485 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

### ZNF485 antibody - N-terminal region - Protein Information

**Name** ZNF485

**Function**
May be involved in transcriptional regulation.

**Cellular Location**
Nucleus.


### ZNF485 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

**ZNF485 antibody - N-terminal region - Images**




WB Suggested Anti-ZNF485 Antibody Titration: 0.2-1 µg/ml
Positive Control: Human Liver

**ZNF485 antibody - N-terminal region - References**

Deloukas,P., (2004) Nature 429 (6990), 375-381 Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.