

ZBTB9 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11649**Specification**

ZBTB9 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O96C00
Other Accession	NM_152735 , NP_689948
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine
Predicted	Rat, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51kDa KDa

ZBTB9 antibody - C-terminal region - Additional Information**Gene ID** 221504**Alias Symbol** MGC23166, ZNF919**Other Names**

Zinc finger and BTB domain-containing protein 9, ZBTB9

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZBTB9 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZBTB9 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZBTB9 antibody - C-terminal region - Protein Information**Name** ZBTB9**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

ZBTB9 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZBTB9 antibody - C-terminal region - Images



WB Suggested Anti-ZBTB9 Antibody Titration: 0.2-1 μ g/ml

Positive Control: Transfected 293T

ZBTB9 antibody - C-terminal region - References

Gerhard,D.S., Genome Res. 14 (10B), 2121-2127 (2004)Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.