

Aebp2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11651**Specification**

Aebp2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O91YQ3
Other Accession	NM_145473 , NP_663448
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa KDa

Aebp2 antibody - C-terminal region - Additional Information**Gene ID** 105859**Alias Symbol** **AU023766, B230313N05Rik****Other Names**

Cold shock domain-containing protein C2, RNA-binding protein PIPPIn, Csdc2, Pippin

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Aebp2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Aebp2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Aebp2 antibody - C-terminal region - Protein Information**Name** Csdc2**Synonyms** Pippin**Function**

RNA-binding factor which binds specifically to the very 3'-UTR ends of both histone H1 and H3.3 mRNAs, encompassing the polyadenylation signal. Might play a central role in the negative regulation of histone variant synthesis in the developing brain (By similarity).

Cellular Location


Nucleus. Cytoplasm. Note=PIPPin-RNA complexes are located to the nucleus.

Aebp2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Aebp2 antibody - C-terminal region - Images



WB Suggested Anti-Aebp2 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Spleen